

A Recurrence Formula for Solutions of Burgers Equations *

Hongling Su¹ Mingliang Wang² Mengzhao Qin¹

¹ CAST (World Laboratory),
Institute of Computational Mathematics
and Scientific/Engineering Computing,
Academy of Mathematics and System Sciences,
Chinese Academy of Sciences, Beijing, 100080, China
²Lanzhou University

February 7, 2008

Abstract

A Bäcklund transformation(BT) and a recurrence formula are derived by the homogeneous balance(HB) method. A initial problem of Burgers equations is reduced to a initial problem of heat equation by the BT, the initial problem of heat equation is resolved by the Fourier transformation method, substituting various solutions of the initial problem of the heat equation will yield solutions of the initial problem of the Burgers equations.

1 Introduction

The HB method is used to find solitary wave solutions and other kinds of exact solutions of nonlinear partial differential equations (PDEs)^[1,2], in this paper, the HB method is used first time with the help of the Fourier transformation method to solve some initial problems of Burgers equations which have been widely used in physical system.

This paper falls into 3 parts. In section 2, by the HB method we obtain a general BT^{3,4} which shows both the connection between solutions of Burgers equations itself and connection between solutions of Burgers and heat equations, its special case is Cole–Hopf transformation and especially

* Supported by the Special Funds for Major State Basic Research Projects , G 1999, 032800

through the BT a recurrence formula is generated for solutions of Burgers equations as

$$\begin{aligned} u_{N+1} &= \frac{u_{Nx} + v_{Nx}}{u_N + v_N} + u_N, \\ v_{N+1} &= \frac{u_{Ny} + v_{Ny}}{u_N + v_N} + v_N. \end{aligned}$$

A reasonable initial problem of the Burgers equations is put out and reduced to a initial problem of the heat equation which is solved by Fourier transformation method in Sec.3. The solution of the initial problem of the heat equation with the aid of the Burgers yields the solution of the initial problem of the Burgers.

2 Burgers Equations

$$\begin{aligned} u_t &= u_{xx} + u_{yy} + 2uu_x + 2vv_y, \\ v_t &= v_{xx} + v_{yy} + 2uv_x + 2vv_y. \end{aligned} \tag{1}$$

This is the famous Burgers equations which have got much attention in recent years. In Ref.[7,8,9], high dimensional and high degree Burgers equations are studied, and a pair of exact solution of Burgers equations(1) is obtained in Ref.[4]. In this section we will present some BTs of Burgers equations(1) through the HB method.

According to the idea of HB method, we seek for its solution of the form:

$$\begin{aligned} u &= \frac{\partial f(\varphi)}{\partial x} + u_0 = f'(\varphi)\varphi_x + u_0, \\ v &= \frac{\partial g(\varphi)}{\partial y} + v_0 = g'(\varphi)\varphi_y + v_0, \end{aligned} \tag{2}$$

here $f = f(\varphi)$ and $g = g(\varphi)$ which are functions of single variable, and $\varphi = \varphi(x, y, t)$ are all to be determined later. (u_0, v_0) are a pair of solutions of (1). By (2), it is easy to deduce that

$$\begin{aligned} u_t &= f''\varphi_x\varphi_t + f'\varphi_{xt} + u_{0t}, \\ u_{xx} &= f''' \varphi_x^3 + 3f''\varphi_x\varphi_{xx} + f'\varphi_{xxx} + u_{0xx}, \\ u_{yy} &= f''' \varphi_x\varphi_y^2 + 2f''\varphi_{xy}\varphi_y + f''\varphi_x\varphi_{yy} + f'\varphi_{xyy} + u_{0yy}, \\ 2uu_x &= 2f'f''\varphi_x^3 + 2f'^2\varphi_x\varphi_{xx} + 2f''\varphi_x^2u_0 + 2f'\varphi_{xx}u_0 + 2f'\varphi_xu_{0x} + 2u_0u_{0x}, \\ 2vu_y &= 2g'f''\varphi_x\varphi_y^2 + 2g'f'\varphi_y\varphi_{xy} + 2f''\varphi_x\varphi_yv_0 + 2f'\varphi_{xy}v_0 + 2g'\varphi_yu_{0y} + 2vu_{0y}, \end{aligned} \tag{3}$$

$$\begin{aligned}
v_t &= g''\varphi_y\varphi_t + g'\varphi_{yt} + v_{0t}, \\
v_{xx} &= g''' \varphi_y\varphi_x^2 + 2g''\varphi_x\varphi_{xy} + g''\varphi_y\varphi_{xx} + g'\varphi_{yxx} + v_{0xx}, \\
v_{yy} &= g''' \varphi_y^3 + 3g''\varphi_y\varphi_{yy} + g'\varphi_{yyy} + v_{0yy}, \\
2uv_y &= 2f'g''\varphi_x^2\varphi_y + 2f'g'\varphi_x\varphi_{xy} + 2g''\varphi_x\varphi_yu_0 + 2g'\varphi_{xy}u_0 + 2f'\varphi_xv_{0x} + 2u_0v_{0x}, \\
2vv_y &= 2g'g''\varphi_y^3 + 2g'^2\varphi_y\varphi_{yy} + 2g''\varphi_y^2v_0 + 2g'\varphi_{yy}v_0 + 2g'\varphi_yv_{0y} + 2v_0v_{0y},
\end{aligned} \tag{4}$$

substituting (3) and (4) into (1), first setting the coefficients of φ_x^3 and φ_y^3 to zero, we obtain ODEs

$$\begin{aligned}
f''' + 2f'f'' &= 0, \\
g''' + 2g'g'' &= 0,
\end{aligned} \tag{5}$$

which have solutions $f = g = \ln \varphi$, thereby it holds that

$$\begin{aligned}
f''' + 2g'f'' &= 0, \quad g''' + 2f'g'' = 0, \\
f'' + f'^2 &= 0, \quad g'' + g'^2 = 0,
\end{aligned} \tag{6}$$

by using (3)–(6), we obtain the expressions:

$$\begin{aligned}
&u_t - u_{xx} - u_{yy} - 2uu_x - 2vu_y \\
&= f''(\varphi_x\varphi_t - \varphi_x\varphi_{xx} - \varphi_x\varphi_{yy} - 2\varphi_x\varphi_yv_0 - 2\varphi_x^2u_0) \\
&+ f'(\varphi_{xt} - \varphi_{xxx} - \varphi_{xyy} - 2\varphi_{xx}u_0 - 2\varphi_xu_{0x} - 2\varphi_{xy}v_0 - 2\varphi_yu_{0y}),
\end{aligned} \tag{7}$$

$$\begin{aligned}
&v_t - v_{xx} - v_{yy} - 2uv_x - 2vv_y \\
&= f''(\varphi_y\varphi_t - \varphi_y\varphi_{yy} - \varphi_y\varphi_{xx} - 2\varphi_x\varphi_yu_0 - 2\varphi_y^2v_0) \\
&+ f'(\varphi_{yt} - \varphi_{yxx} - \varphi_{yyy} - 2\varphi_{xy}u_0 - 2\varphi_xv_{0x} - 2\varphi_yv_{0y} - 2\varphi_{yy}v_0),
\end{aligned} \tag{8}$$

It is easy to see that (2) are solutions of (1), provide that the right hand sides of (7) and (8) to be zero, hence set

$$\varphi_t - \varphi_{xx} - \varphi_{yy} - 2\varphi_xu_0 - 2\varphi_yv_0 = 0, \tag{9}$$

$$\varphi_{xt} - \varphi_{xxx} - \varphi_{xyy} - 2\varphi_{xx}u_0 - 2\varphi_xu_{0x} - 2\varphi_{xy}v_0 - 2\varphi_yu_{0y} = 0, \tag{10}$$

$$\varphi_{yt} - \varphi_{yxx} - \varphi_{yyy} - 2\varphi_{yy}v_0 - 2\varphi_yv_{0y} - 2\varphi_{xy}u_0 - 2\varphi_xv_{0x} = 0, \tag{11}$$

then the right hand sides of (7) and (8) respectively vanish. In fact (10) and (11) can be reduced into

$$(\varphi_t - \varphi_{xx} - \varphi_{yy} - 2\varphi_xu_0 - 2\varphi_yv_0)_x + 2\varphi_yv_{0x} - 2\varphi_yu_{0y} = 0, \tag{12}$$

$$(\varphi_t - \varphi_{xx} - \varphi_{yy} - 2\varphi_xu_0 - 2\varphi_yv_0)_y + 2\varphi_xu_{0y} - 2\varphi_xv_{0x} = 0, \tag{13}$$

thereby, (9)–(11) equal that

$$\begin{aligned}\varphi_t - \varphi_{xx} - \varphi_{yy} - 2\varphi_x u_0 - 2\varphi_y v_0 &= 0, \\ u_{0y} - v_{0x} &= 0,\end{aligned}\tag{14}$$

here $\varphi_x^2 + \varphi_y^2 \neq 0$. Substituting $f = g = \ln \varphi$ into (2), we obtain BT of Burgers equations (1)

$$u = \frac{\varphi_x}{\varphi} + u_0,\tag{15}$$

$$v = \frac{\varphi_y}{\varphi} + v_0,\tag{16}$$

$$\varphi_t - \varphi_{xx} - \varphi_{yy} - 2\varphi_x u_0 - 2\varphi_y v_0,\tag{17}$$

$$u_{0y} - v_{0x} = 0,\tag{18}$$

namely, (u_0, v_0) is a pair of solutions of Burgers equations (1), and $u_{0y} = v_{0x}$ therefore (2) is another pair of solutions of (1) only if $\varphi = \varphi(x, y, t)$ satisfies linear equation (9). Now set $\varphi = u_0 + v_0$, (9) is automatically satisfied, additionally, if $u_{0y} = v_{0x}$ exists in the first pair of solutions (u_0, v_0) , the new solutions (u, v) according to the BT have to satisfy $u_y = v_x$, the circular use of this *BT* yields the recurrence formula of Burgers equations (1) as

$$\begin{aligned}u_{N+1} &= \frac{u_{Nx} + v_{Nx}}{u_N + v_N} + u_N, \\ v_{N+1} &= \frac{u_{Ny} + v_{Ny}}{u_N + v_N} + v_N.\end{aligned}\tag{19}$$

Here $N = 0, 1, 2, \dots$, with $v_{0x} = u_{0y}$.

In addition, if we take initial solution of Burgers equations (1) as $u_0 = v_0 = 0$ (the trivial solutions of Burgers equations (1)), BT (15)–(18) respectively reduce to

$$\begin{aligned}u &= \frac{\varphi_x}{\varphi}, & v &= \frac{\varphi_y}{\varphi}, \\ \varphi_t - \varphi_{xx} - \varphi_{yy} &= 0,\end{aligned}\tag{20}$$

which is usually called Cole–Hopf transformation.

3 Fourier Transformation

Provided the initial value problem of Burgers equations in what follows

$$u_t = u_{xx} + u_{yy} + 2uu_x + 2vu_y,\tag{21}$$

$$v_t = v_{xx} + v_{yy} + 2uv_x + 2vv_y,\tag{22}$$

$$u|_{t=t_0} = s(x, y), \quad v|_{t=t_0} = k(x, y)\tag{23}$$

where $s(x, y), k(x, y) \rightarrow 0$ when $|x|, |y| \rightarrow 0$.

By using the former result, we can take its solution to be as

$$u = \frac{\varphi_x}{\varphi} + u_0, \quad v = \frac{\varphi_y}{\varphi} + v_0, \quad (24)$$

here (u_0, v_0) are constant solutions of (21)–(22) and φ satisfies

$$\varphi_t - \varphi_{xx} - \varphi_{yy} - 2u_0\varphi_x - 2v_0\varphi_y = 0 \quad (25)$$

then (21)–(23) can be rewritten into the initial value problem of the following form

$$\begin{aligned} \varphi_t - \varphi_{xx} - \varphi_{yy} - 2u_0\varphi_x - 2v_0\varphi_y &= 0, \\ \varphi|_{t=t_0} &= f(x, y), \end{aligned} \quad (26)$$

which

$$f(x, y) = e^{\int_{x_0}^x s_1(x, y_0) dx + \int_{y_0}^y k_1(x, y) dy} \quad (27)$$

(x_0, y_0) are arbitrary constants, $s_1(x, y) = s(x, y) - u_0$, $k_1(x, y) = k(x, y) - v_0$. Then if u_0, v_0 are arbitrary constant solutions of equations (1), by using of Fourier transformation, we obtain solutions of (26)–(27).

Now introduce the Fourier transformation simply. If $f(x, y, t)$ is a function, $\widehat{f}(c_1, c_2, t)$ is used to denote its Fourier transformation as follow:

$$\widehat{f}(c_1, c_2, t) = F[f(x, y, t)] = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y, t) e^{-ic_1 x - ic_2 y} dx dy \quad (28)$$

and

$$f(x, y, t) = F^{-1}[\widehat{f}(c_1, c_2, t)] = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \widehat{f}(c_1, c_2, t) e^{ic_1 x + ic_2 y} dc_1 dc_2 \quad (29)$$

we'd like show its several properties which are used here:

- I $F[af + by] = aF[f] + bF[g]$
- II $F[\frac{\partial^m \partial^n f}{\partial x^m \partial y^n}] = (ic_1)^m (ic_2)^n F[f]$
- III $F[f(x - a, y - b, t)] = e^{-ic_1 a - ic_2 b} F[f(x, y, t)]$
- IV $F[f(ax, by, t)] = \frac{1}{|ab|} \widehat{f}(\frac{c_1}{a}, \frac{c_2}{b}, t)$
- V if $(f * g)(x, y, t) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x - z_1, y - z_2, t) g(z_1, z_2, t) dz_1 dz_2$, then

$$F^{-1}[\widehat{f\widehat{g}}] = \frac{1}{2\pi} f * g$$

here, a and b are arbitrary constants, m and n are arbitrary integers. In order to deal with the problem (26), the Fourier transformation is done for them, it is reduced to a order differential problem as following:

$$\begin{aligned} \frac{d\widehat{\varphi}}{dt} + c_1^2\widehat{\varphi} + c_2^2\widehat{\varphi} - 2iu_0c_1\widehat{\varphi} - 2v_0c_2\widehat{\varphi} &= 0, \\ \widehat{\varphi}|_{t=t_0} &= \widehat{f}(x, y), \end{aligned} \quad (30)$$

which admits solutions as

$$\widehat{\varphi}(c_1, c_2, t) = \widehat{f}(c_1, c_2, t) e^{-(c_1^2 + c_2^2 - 2iu_0c_1 - 2iv_0c_2)(t-t_0)} \quad (31)$$

set $\widehat{g} = e^{-(c_1^2 + c_2^2 - 2iu_0c_1 - 2iv_0c_2)(t-t_0)}$ such that

$$g(x, y, t) = F^{-1}[\widehat{g}(c_1, c_2, t)] = \frac{1}{2(t-t_0)} e^{-\frac{(x-2u_0)^2 + (y-2v_0)^2}{4(t-t_0)}} \quad (32)$$

so the solution of (26) is gained

$$\begin{aligned} \varphi(x, y, t) = F^{-1}[\widehat{f}\widehat{g}] &= \frac{1}{4\pi(t-t_0)} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{\int_{x_0}^{x-z_1} s_1(\tau_1, y_0) d\tau_1 + \int_{y_0}^{y-z_2} k_1(x-z_1, \tau_2) d\tau_2} \\ &\quad e^{-\frac{(z_1-2iu_0)^2 + (z_2-2iv_0)^2}{4(t-t_0)}} dz_1 dz_2, \end{aligned} \quad (33)$$

hence $u = \frac{\varphi_x}{\varphi} + u_0$, $v = \frac{\varphi_y}{\varphi} + v_0$ which are solutions of (21)–(23) also can be got, detail discussion does not present here.

References

- [1] M. L. Wang, Yubin Zhou and Zhibin Li, Applications of a homogeneous balance method to exact solutions of nonlinear equations in mathematical physics, Physics Letters A 216(1996), 67-75.
- [2] M. L. Wang and Z. B. Li, Applications of Quasi-solution Technique to the Exact Solutions of Some Nonlinear Partial Differential Equations, Proceedings of the 1994 Beijing Symposium on Nonlinear Evolution Equations and Infinite Dimensional Dynamical Systems., Zhongshan University Press (1995).
- [3] C. Rogers and W. F. Shodwich, Bäcklund transformations and their applications, Academic Press, Inc, New York, 1982.

- [4] Lamb G. L. Jr, Bäcklund transformation for certain nonlinear evolution equations, J.Math phys, 15(1974), 2157-2165.
- [5] Airy. G. B., Trans. Cambridge Philos. Soc., 6(1838), 379-402
- [6] Landaus. L. D., and Lifshitz. E. M., the classical theory of fields, Addison-Wesley, 1951.
- [7] K-M. Tamizhmani and Punithavathi, Similarity and Painleve property of the coupled higher dimensional Burgers equations, Int.nonlinear mechanics 3 4(1991)429-438.
- [8] Y.Murakami, J. Phys. Soc. Japan, 59(1990), 1.
- [9] M. L. Wang, Acta Mathematica Scientia 6(1986), 335.
- [10] M. Boiti, J. J. P. Leon, M. Manna, and F.Pempinelli, Inv. Problems 2, 271(1986).
- [11] R. Radha and e, J. Math Phys, 35(1994), 4764.
- [12] K. W. Chow, J. Phys. Soc. Jun, 65(1996), 1971.